

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Causes moderate eye irritation. Harmful if swallowed. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear long-sleeved shirt and long pants, socks, shoes and chemical-resistant gloves.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical-resistant to this product are listed below. Mixers, loaders, applicators and other handlers, (including during mixing/loading, repair and cleaning of equipment and disposal of pesticide) must wear the following:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material
- Shoes plus socks

In addition to the above PPE, applicators using high-pressure handwand in an enclosed area must wear at least a NIOSH-approved respirator with a dust/mist filter with MSHA/NIOSH approval prefix TC-21C or any N, R, P, or HE filter.

Applicators using hand-held foggers or fog/mist generators in indoor spaces must wear coveralls over long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material, chemical-resistant footwear plus socks, chemical-resistant headgear, if overhead exposure, and a half-face, full-face or hood style NIOSH-approved respirator with a dust/mist filtering cartridge (MSHA/NIOSH approval number prefix TC-21C), or a cartridge or canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G), or a cartridge or canister with any N, R, P, or HE filter.

In addition to the above PPE, mixers, loaders, applicators and other persons exposed to this product's concentrate, or handlers performing animal dip applications, must wear a chemical-resistant apron.

**User Safety Recommendations:**

- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.
- Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates. Do not apply directly to or near water, or to areas where surface water is present or to intertidal areas below the mean water mark. Do not apply when weather conditions favor drift from treated areas. Drift and run-off from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters. See Directions for Use for additional precautions and use restrictions.

Under some conditions, this product may also have a potential for transport into surface water in run-off (primarily adsorbed to suspended soil particles), for several months or more after application. These include poorly draining or wet soils with readily visible slopes towards adjacent surface waters, frequently flooded areas and areas overlying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips and areas over-lying tile drainage systems that drain to surface waters.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

**PHYSICAL OR CHEMICAL HAZARDS**

Do not use this product in or on electrical equipment due to possibility of shock hazard.

**DIRECTIONS FOR USE**

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

**READ THIS LABEL IN ITS ENTIRETY BEFORE USING.**

**FOR PET USE – USE ONLY ON DOGS OR PUPPIES OVER 12 WEEKS OF AGE**

**USE RESTRICTIONS**

- Apply this product only as specified on this label.
- Do not apply to puppies less than 12 weeks old.
- **DO NOT USE ON CATS OR KITTENS.** Do not treat cats or their bedding. This product may be toxic or potentially fatal if applied to or ingested by cats or kittens.
- Application with aerial or motorized ground equipment is prohibited.
- Do not use spoons or measuring cups for food purposes after use with pesticides.
- Do not use in food areas of restaurants or food handling establishments or other areas where food is commercially prepared or processed. Do not make space spray applications in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms, but excluding areas where foods may be prepared or held. When used in the home, cover or remove all exposed food and cooking utensils and cover all food-processing surfaces or wash thoroughly after treatment and before re-use.
- Do not contaminate food or feedstuffs.
- Except when applying to pets or livestock, do not apply this product in a way that will contact workers or other persons, either directly or through drift. Do not breathe vapors or spray mist.
- Except when applying to pets or livestock, do not enter or allow others to enter until vapors, mists and aerosols have dispersed and the treated area has been thoroughly ventilated.
- Except when applying to pets or livestock, do not apply this product in a way that will contact adults, children or pets, either directly or through drift.
- Except when applying to pets or livestock, only protected handlers may be in the area during application.
- Do not remain in treated area. Exit area immediately and remain outside the treated area until vapors, mists and aerosols have dispersed.
- Remove or cover exposed food and drinking water before application.
- Do not use in aircraft cabins.
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers or storm drains, water bodies, or aquatic habitat can occur. Do not allow this product to enter any drain during or after application.
- Do not make applications during rain.
- Do not wet plants to the point of run-off or drip.
- Do not wet articles to the point of run-off or drip. Do not use treated article until spray has dried. Dry clean treated clothing before wearing.
- When used in dairy barns or facilities: Close milk bulk tank lids to prevent contamination from dead or falling insects. Remove or cover milking utensils before application. Wash teats of animals before milking.

- For direct application to non-domestic animals/livestock: Do not apply more than 1 time per day.
- All outdoor applications must be limited to spot or crack-and-crevice treatments only except for the following permitted uses: (1) Treatment to soil or vegetation around structures; (2) Applications to lawns, turf and other vegetation; (3) Applications to building foundations up to a maximum height of 3 feet. Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors and eaves) are limited to spot and crack-and-crevice applications only.
- Remove pets, birds and cover fish aquariums before spraying. Turn off aquarium filtering systems before spraying and wait approximately 15 minutes after application before turning the filter back on.
- This product is not for use on plants being grown for sale or other commercial use. This product should not be used for research purposes, for commercial seed production, for commercial nursery use or forestry use.

When diluting, add the correct amount of *Pyranha Stock Guard® Concentrate* to water as directed and blend thoroughly. For use both indoors (in homes) and outdoors. May be used in nonfood areas of commercial buildings, warehouses, food/feed processing establishments, institutions, schools, office buildings, hotels, motels, restaurants, theaters, hospitals, horse barns, swine and poultry houses, milking parlors, dairies, cattle barns and kennels. Examples of non-food areas are: lavatories, offices, locker rooms, garbage rooms, entries and vestibules, machine rooms, boiler rooms, garages, mop closets and storage rooms/closets (after canning or bottling).

For maximum effectiveness, a combination of space treatment and localized application is recommended.

**Note:** In the chart below, the number in the 3<sup>rd</sup> column refers to the level of Permethrin in the final diluted product. The rates on this label make the ready-to-use products listed below.

DILUTION RATE	OUNCES OF PYRANHA STOCK GUARD® CONCENTRATE/GALLON	% PERMETHRIN IN DILUTED PRODUCT
1:19	6.40	0.50%
1:39	3.20	0.25%
1:100	1.28	0.10%
1:200	0.64	0.05%
1:400	0.32	0.025%

**WHEN USING INDOORS:**

**SURFACE SPRAY APPLICATION METHODS:**

Insects Controlled	Directions for Use	Dilution Rates
Cockroaches, Waterbugs, Palmetto Bugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Millipedes, Centipedes, Sowbugs, Pillbugs, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Saw-toothed Grain Beetles, Spider Beetles, Cigarette Beetles, Drug Store Beetles, Darkling Beetles, Mealworms, Grain Mites, Cadelles, and Lesser Houseflies and other manure breeding flies.	Adjust your sprayer to deliver a coarse wet spray. Direct the spray into cracks and crevices, insect hiding places, behind shelves and drawers, under pallets, around containers of stored foods, and around the base of machinery. Spray ant trails, nests and points of entry. Spray bookcases for silverfish. Only when used as a surface application, spray floors, walls and other surfaces.	For initial clean up of severe infestation, dilute at a rate of 1 part <i>Pyranha Stock Guard® Concentrate</i> in 19 parts water (6.4 fluid ounces per gallon). This makes a 0.5% permethrin ready-to-use product.  For normal infestation, dilute 1 part <i>Pyranha Stock Guard® Concentrate</i> in 39 parts water (3.2 ounces per gallon).  For maintenance use, dilute 1 part <i>Pyranha Stock Guard® Concentrate</i> in 100 parts water (1.28 ounces/gallon or 8 teaspoons/gallon). This makes a 0.1% permethrin ready-to-use product.
Centipedes, Ground Beetles, Pillbugs, Sowbugs, Scorpions and Ticks	Apply around windows and doors as well as other places where these insect pests may enter premises. Treat storage areas, baseboards, and other places where these pests may hide.	For maintenance use, dilute 1 part <i>Pyranha Stock Guard® Concentrate</i> in 100 parts water (1.28 ounces/gallon or 8 teaspoons/gallon). This makes a 0.1% permethrin ready-to-use product.
Carpenter Beetles	Spray edges of and under carpeting and rugs. Make localized application to baseboards and floors. Spray directly into cracks and crevices, closets and infested areas of shelving.	
Flying Insect places where they rest or land (helps in controlling insects such as Gnats, Mosquitoes, Skippers, Boxelder Bugs, Asian Lady Beetles)	Treat walls, ceilings, moldings, screens, door and window frames and similar resting places.	Dilute at a rate of 1 part <i>Pyranha Stock Guard® Concentrate</i> in 19 parts water (6.4 fluid ounces per gallon). This makes 0.5% permethrin ready-to-use product.

**SURFACE SPRAY APPLICATION METHODS:**

Bed Bugs	Before treating, spray fabric in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying. If no staining occurs, continue by spraying mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings, and floors. Allow all treated articles to thoroughly dry before use.	For initial clean up of severe insect infestation, dilute at a rate of 1 part <i>Pyranha Stock Guard® Concentrate</i> in 19 parts water (6.4 fluid ounces per gallon). Makes a 0.5% permethrin ready-to-use product.
Clothes Moths	Remove infested items from storage, brush thoroughly and if possible, air for a few hours, preferably in sunlight. Liberally apply spray to empty closets, chests, dressers and other storage areas by directing the spray into cracks, crevices and joints. After airing, infested items may be treated lightly, keeping nozzle at least 3 feet from fabric to avoid staining.	For normal infestations, dilute 1 part <i>Pyranha Stock Guard® Concentrate</i> in 39 parts water (3.2 ounces per gallon). Will make a 0.25% permethrin ready-to-use product.
Lice and Louse Eggs (on inanimate objects)	Prior to using, spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry-cleaned.	
Fleas, Flies, Ticks, Lice and Mange Mites (adults and larvae) in kennels, shelters, doghouses, runs and yards	Allow all treated articles to dry thoroughly before use. Test delicate fabrics in an inconspicuous area prior to use. Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames and localized areas of floor and floor coverings (rugs and carpets). Fresh bedding should be placed in animal quarters once spray has dried. Adult fleas and larvae contacted by spray will be killed. Do not treat cat bedding.	For maintenance use, dilute 1 part <i>Pyranha Stock Guard® Concentrate</i> in 100 parts water (1.28 ounces per gallon or 8 teaspoons per gallon). Makes a 0.1% permethrin ready-to-use product.

**SPACE SPRAY APPLICATION METHODS:**

Insects Controlled	Directions for Use	Dilution Rates
House Flies, Fruit Flies, Gnats, Mosquitoes, Skipper Flies, Wasps, Hornets, Bees, Black Flies, Asian Lady Beetles, Boxelder Bugs, Angoumois Grain Moths, and Tobacco Moths	Apply with non thermal mechanical or compressed air equipment that is adjusted to deliver a fine mist. Close doors and windows and shut off ventilating systems, including air conditioners. Direct spray at an upward angle distributing it uniformly through the entire area at a rate of 1 ounce per 1,000 cubic feet of space. Vacate after treatment and keep area closed for at least 10 minutes then ventilate and sweep up dead insects before occupying. For quick kill of accessible or exposed stages of other insects named on this label. Apply using conventional non thermal mechanical or compressed air equipment. Follow space spraying directions. Apply dilutions at a rate of 1 ounce per 1,000 cubic feet of space.	For initial clean up of severe insect infestation, dilute at a rate of 1 part <i>Pyranha Stock Guard® Concentrate</i> in 19 parts water (6.4 fluid ounces per gallon). Makes a 0.5% permethrin ready-to-use product.  For normal infestations, dilute 1 part <i>Pyranha Stock Guard® Concentrate</i> in 39 parts water (3.2 ounces per gallon). Will make a 0.25% permethrin ready-to-use product.  For maintenance use, dilute 1 part <i>Pyranha Stock Guard® Concentrate</i> in 100 parts water (1.28 ounces per gallon or 8 teaspoons/gallon). Makes a 0.1% permethrin ready-to-use product.

(Continued)

### OUTDOOR USE AREAS

Drive-In Theaters, Drive-In Restaurants, Gardens, Homes, Parks, Playgrounds, Recreational Areas, Golf Courses, Feed Lots, Corrals, Swine Lots, Zoos and Urban Areas.

### PREMISE SPRAY APPLICATION METHODS:

Insects Controlled	Directions for Use	Dilution Rates
Mosquitoes, Wasps, Hornets, Yellow Jackets, Annoyance Flies and Skippers	Treat walls, ceilings, moldings, screens, door and window frames, and similar resting places. Application should be made prior to occupancy. Cover any exposed food, water, drinking fountains and animal feed before application. In zoos, prevent reptiles from exposure to this product.	Dilute 1 part <i>Pyranha Stock Guard® Concentrate</i> in 39 parts water (3.2 fluid ounces per gallon of water). This will make a 0.25% permethrin ready-to-use product.
Gypsy Moth Caterpillars and Adults, Adult Japanese Beetles, Eastern and Forest Tent Caterpillars, Mole Crickets and Grasshoppers	Apply to insure thorough coverage of upper and lower leaf surfaces. Thoroughly wet down grass a few hours before applying. Lawns should not be higher than 3 inches at time of treatment. Wet lawns with 1/4 inch of water immediately after application. For heavy infestations, repeat application after 2 weeks.  Treat trees and shrubs as soon as insects emerge (usually late spring). Spray leaves, branches, and tree trunks evenly. Make sure to penetrate dense foliage. Repeat application every 7 days as necessary for control.  On roses and in flower gardens: Treat plants and flowers as soon as insects appear. Spray flowers and plants making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.	Dilute 1 part concentrate in 39 parts water (3.2 fluid ounces per gallon of water). Makes a 0.25% permethrin ready-to-use product.  Mix 1.28 fl. oz. per gallon of water and treat 1,000 sq. ft. Slightly more water may be used provided the amount of product per area is the same.
Fire Ants	Using equipment such as a garden sprinkling can, gently apply at the rate of one gallon per mound. Thoroughly wet the mound and surrounding area to a diameter of 4 feet. For best results, apply in cool weather, 65° to 80°F, or in early morning or late evening hours. Treat new mounds as they appear. Do not disturb the mound prior to treatment. Treat all mounds in the vicinity and treat all colonies, which have not yet built mounds. Do not use equipment that produces a pressurized spray since this will disturb Fire Ants and cause migration, reducing the effectiveness of this product.	Mix 6.4 fl. oz. in 1 gallon of water.

8

### PREMISE SPRAY APPLICATION METHODS:

(Continued)

Ants, Crickets, Earwigs, Fleas and Ticks	For localized infestations where there are weeds or bushy non-crop areas, spray infested areas thoroughly. For ants, thoroughly wet hills and runways.	For severe infestation, dilute 1 part <i>Pyranha Stock Guard® Concentrate</i> in 19 parts water (6.4 oz. per gallon of water). Will make a 0.5% permethrin ready-to-use product. <b>Also</b> treat 400 linear feet with a 6-inch spray band. For normal infestations, dilute 1 part <i>Pyranha Stock Guard® Concentrate</i> in 39 parts water (3.2 oz. per gallon). Makes a 0.25% permethrin ready-to-use product.
Clover Mites, Millipedes, Sowbugs (Pillbugs)	To help prevent infestation of buildings, treat the building foundation to a height of 2 to 3 feet where insects are active and may find entry. Apply also as a residual spray to outside surfaces of buildings, garages, patios, and other areas where these pests may be found.	
Armyworms, Cutworms, Chinch Bugs, Sod Webworms	Thoroughly wet down grass a few hours before applying. Lawns should be no higher than 3 inches at time of application. Delay watering or mowing for 24 hours after treatment. For heavy infestation, repeat application after 2 weeks.	Mix 3.2 fluid ounces per gallon of water and treat 1,000 sq. ft. Slightly more water may be used as long as the amount of product per area is the same.

### ANIMALS AND ANIMAL PREMISE USE AREAS

Cattle Barns, Milk Rooms & Parlors, Dairies, Horse Barns, Poultry Houses, Swine Houses, and Kennels.

This product may be used on Dairy and Beef Cattle, Sheep, Goats, Swine, Horses, Poultry and Dogs.

Insects Controlled	Directions for Use	Dilution Rates
Horn Flies, House Flies, Mosquitoes and Gnats	Wet the hair thoroughly, paying particular attention to topline, underline, flanks, withers and other infested areas. Repeat treatment at intervals of 5 to 12 days for small insect populations or as needed when flies are emerging in large numbers.	For clean up of severe insect infestation, dilute at a rate of 1 part <i>Pyranha Stock Guard® Concentrate</i> in 100 parts water (1.28 ounces per gallon or 8 teaspoons/gallon). Makes a 0.1% permethrin ready-to-use product.
Stable Flies, Horse Flies, and Deer Flies	Apply a quart per adult animal to wet the hair thoroughly with special attention to the legs, flanks, barrel, topline and other body areas commonly attacked by those flies. Repeat daily as needed.	For normal infestations, dilute 1 part <i>Pyranha Stock Guard® Concentrate</i> in 200 parts water (0.64 ounces per gallon or 4 teaspoons/gallon). Makes a 0.05% permethrin ready-to-use product.
Face flies	Apply using spray that produces large wetting droplets. Apply to the face of the animal in the morning before moving it to pasture. Apply sufficiently to wet the face, but not more than 1.5 ounces per animal. Do not get into eyes, nose or mouth. Repeat daily as needed.	For maintenance use, dilute 1 part <i>Pyranha Stock Guard® Concentrate</i> in 400 parts water (0.32 ounces per gallon or 2 teaspoons per gallon). Makes a 0.025% permethrin ready-to-use product.
Poultry Lice	Spray roosts, nests, walls and cages thoroughly, followed by spraying over the birds with a fine mist.	

9

### ANIMALS AND ANIMAL PREMISE USE AREAS

(Continued)

Insects Controlled	Directions for Use	Dilution Rates
Bed Bugs and Mites (including Northern fowl mite) in poultry houses	Spray crevices of roost poles, cracks in nests and walls where the Mites & Bed Bugs hide. Spray birds with a fine mist paying particular attention to vents. Use at the rate of 1 gallon of spray per 100 birds.	For cleanup of severe insect infestation, dilute at a rate of 1 part <i>Pyranha Stock Guard® Concentrate</i> in 100 parts water (1.28 ounces per gallon or 8 teaspoons/gallon). Makes a 0.1% permethrin ready-to-use product.
Sheep "Tick" or Ked	Thoroughly wet all parts of the animal's body by dipping or spraying with sufficient pressure to penetrate the wool. Treat at a rate sufficient to wet the animal.	For normal infestations, dilute 1 part <i>Pyranha Stock Guard® Concentrate</i> in 200 parts water (0.64 ounces per gallon or 4 teaspoons per gallon). Makes a 0.05% permethrin ready-to-use product.
Meal Worms, Darkling Beetles	Apply as a fine mist or fog, directing the spray toward the upper corners and ceiling until the area is filled with mist. Use about 2 ounces per 1,000 cubic feet of space. For best results, close windows and doors before spraying. Vacate after treatment and keep area closed for 10 to 15 minutes. Ventilate area before reentering the area.	For maintenance use, dilute 1 part <i>Pyranha Stock Guard® Concentrate</i> in 400 parts water (0.32 ounces per gallon or 2 teaspoons per gallon). Makes a 0.025% permethrin ready-to-use product.
Horn Flies, Face Flies, Horseflies, Ear Ticks	Back rubber – make sure the back rubber stays filled.	Dilute 8 oz. in 5 gallons oil.
Ticks (including deer ticks, which carry Lyme disease) Blowflies, Hog Lice	Spray, dip, or sponge to apply 1 pint per animal, especially around ears.	Dilute 1.28 oz. per gallon of water (1 part <i>Pyranha Stock Guard® Concentrate</i> to 100 parts water). This makes a 0.10% permethrin ready-to-use product.
Poultry Mites, Northern Fowl Mites and Lice	At the rate of 1 gallon per 100 birds, spray with a fine mist. Spray walls, roosts, cages and nests.	
Fleas, Flies, Ticks, Lice and Mange Mites (adults and larvae) in doghouses, kennels, shelters, yards and runs.	Before treating, spray fabric in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying. If no staining occurs, thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and doorframes and localized areas of floor and floor coverings (rugs and carpets). Fresh bedding should be placed in animal quarters once spray has dried. Adult fleas and larvae contacted by spray will be killed.  Do not treat cat bedding.	For initial clean up of severe insect infestation, dilute at a rate of 1 part <i>Pyranha Stock Guard® Concentrate</i> in 19 parts water (6.4 fluid ounces/gallon). This makes a 0.5% permethrin ready-to-use product.  For normal infestations, dilute 1 part <i>Pyranha Stock Guard® Concentrate</i> in 39 parts water (3.2 ounces per gallon). Makes a 0.25% permethrin ready-to-use product.  For maintenance use, dilute 1 part concentrate in 100 parts water (1.28 ounces/gallon or 8 teaspoons/gallon). Makes a 0.1% permethrin ready-to-use product.

10

### ANIMALS AND ANIMAL PREMISE USE AREAS

(Continued)

Fleas, Ticks, Lice and Mange Mites (adults and larvae) on dogs and livestock	Thoroughly wet the animal by spraying, dipping or sponging. Do not rinse or towel dry; allow animal to air dry in a warm place. Gives long lasting protection against reinfection. Repeat applications at intervals of 2 to 3 weeks.  <b>Do not treat cats or their bedding. Do not treat puppies less than twelve weeks old.</b>  Pet sensitivities may occur after using ANY pesticide product. If you see signs of sensitivity bathe your pet with mild soap and rinse with large quantities of water. If signs continue, consult a veterinarian immediately.  If you have a debilitated, aged, medicated, pregnant or nursing animal, consult a veterinarian before using this product.	For clean up of severe insect infestation, dilute at a rate of 1 part concentrate in 100 parts water (1.28 ounces per gallon or 8 teaspoons/gallon). Makes a 0.1% permethrin ready-to-use product.  For normal infestations, dilute 1 part concentrate in 200 parts water (0.64 ounces per gallon or 4 teaspoons/gallon). Makes a 0.05% permethrin ready-to-use product.
--	---	--

FOR DAIRY CATTLE: Spray lactating dairy cows only after milking is complete.

FOR SWINE: Do not ship animals for slaughter within 5 days of last treatment.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store upright at room temperature. Avoid exposure to extreme temperatures.

**CONTAINER DISPOSAL:** Non-refillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

11

### ANIMALS AND ANIMAL PREMISE USE AREAS

(Continued)

Fleas, Ticks, Lice and Mange Mites (adults and larvae) on dogs and livestock	Thoroughly wet the animal by spraying, dipping or sponging. Do not rinse or towel dry; allow animal to air dry in a warm place. Gives long lasting protection against reinfection. Repeat applications at intervals of 2 to 3 weeks.  <b>Do not treat cats or their bedding. Do not treat puppies less than twelve weeks old.</b>  Pet sensitivities may occur after using ANY pesticide product. If you see signs of sensitivity bathe your pet with mild soap and rinse with large quantities of water. If signs continue, consult a veterinarian immediately.  If you have a debilitated, aged, medicated, pregnant or nursing animal, consult a veterinarian before using this product.	For clean up of severe insect infestation, dilute at a rate of 1 part concentrate in 100 parts water (1.28 ounces per gallon or 8 teaspoons/gallon). Makes a 0.1% permethrin ready-to-use product.  For normal infestations, dilute 1 part concentrate in 200 parts water (0.64 ounces per gallon or 4 teaspoons/gallon). Makes a 0.05% permethrin ready-to-use product.
--	---	--

FOR DAIRY CATTLE: Spray lactating dairy cows only after milking is complete.

FOR SWINE: Do not ship animals for slaughter within 5 days of last treatment.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store upright at room temperature. Avoid exposure to extreme temperatures.

**CONTAINER DISPOSAL:** Non-refillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

PYRANHA is a registered trademark of PYRANHA, INC.  
All rights reserved  
REV103116

12

**Pyranha®**

## PYRANHA STOCK GUARD® CONCENTRATE

### FIRST AID

<b>If in Eyes:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses if present, after the first 5 minutes, then continue rinsing.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If Swallowed:</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on Skin or Clothing:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>

### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For additional information on this product (including health concerns, medical emergencies or pesticide incidents), you may call 1-800-222-1222, twenty-four (24) hours per day, seven (7) days per week.

EPA Reg. No. 1021-2767-21165

EPA Est. No. 42403-TX-001

Manufactured for  
**PYRANHA, INC.**  
6602 CUNNINGHAM ROAD  
HOUSTON, TX 77041  
800-231-2966



9 1738 11464 9

1

Peel Here →