



NDC 17135-173-03

# PyrantelCare™ Daily Dewormer 2.11%

(pyrantel tartrate)

## Equine Anthelmintic

### Medicated for Continuous Feeding

For the prevention of *Strongylus vulgaris* larval infestation in horses

For control of the following parasites in horses:

**LARGE STRONGYLES** (adults) - *S. vulgaris*, *S. edentatus*

**SMALL STRONGYLES** (adults and fourth-stage larvae) - *Cyathostomum* spp.,  
*Cylicocycclus* spp., *Cylicostephanus* spp., *Cylicodontophorus* spp.,  
*Poteriostomum* spp., *Triodontophorus* spp.

**PINWORMS** (adults and fourth-stage larvae) - *Oxyuris equi*

**ASCARIDS** (adults and fourth-stage larvae) - *Parascaris equorum*

#### ACTIVE DRUG INGREDIENT:

Pyrantel tartrate ..... 2.11% (9.6 g/lb)

#### GUARANTEED ANALYSIS:

Crude Protein (not less than)..... 20.10%

Crude Fat (not less than) ..... 2.56%

Crude Fiber (not more than)..... 18.16%

Moisture (not more than) ..... 7.35%

**INGREDIENTS:** Brewers dried yeast, brewers dried grains, dehydrated alfalfa meal, wheat middlings, natural and artificial flavors, propionic acid (preservative)



Approved by FDA under ANADA # 200-168

(SEE BACK PANEL FOR DIRECTIONS)

**NET WEIGHT: 50 lb (22.6 kg)**

Item: 100521750  
RM: 99.0002189.00  
Rev 03-2019



# PyrantelCare™ Daily Dewormer 2.11%

(pyrantel tartrate)

## Equine Anthelmintic

**MIXING AND FEEDING DIRECTIONS:** PyrantelCare™ Daily Dewormer 2.11% (pyrantel tartrate) is to be administered on a continuous basis either as a top-dress or mixed in the horse's daily grain ration at the rate of 1.2 mg pyrantel tartrate per pound of body weight daily. To achieve this dose, administer 0.5 oz. (½ scoop) of PyrantelCare™ dewormer 2.11% per 250 lb. of body weight. (For example, a 1,000 lb. horse requires 2 oz. or two scoops. See Top-Dress Directions.) Please note that the measuring cup is different in size and shape from the PyrantelCare™ Daily Dewormer 1.06% measuring cup due to the higher concentration of PyrantelCare™ Daily Dewormer 2.11%. PyrantelCare™ dewormer 2.11% should be administered for the entire period that the animal is at risk to internal parasites. Unprotected animals that have grazed may already have an established *S. vulgaris* larval infestation. Before administering PyrantelCare™ dewormer 2.11%, these animals should be treated with a therapeutic dose of a larvicidal product. Foals may be administered PyrantelCare™ dewormer 2.11% as soon as consistent intake of grain mix is occurring. This is generally between 2 to 3 months of age.

PyrantelCare™ dewormer 2.11% may be used in mares at any stage of pregnancy or lactation. Stallion fertility is not affected by the use of PyrantelCare™ dewormer 2.11%.

#### MEDICATED GRAIN MIX DIRECTIONS

LB OF MEDICATED GRAIN MIX PER 100 LB OF BODY WEIGHT	LB OF PYRANTELCARE™ DAILY DEWORMER 2.11%	LB OF NON-MEDICATED FEED	CONCENTRATION GRAMS PER TON
2.0	12.5	1987.5	120
1.5	16.5	1983.5	160
1.0	25.0	1975	240
0.5	50.0	1950	480
0.2	125.0	1875	1200

**CAUTION:** Consult your veterinarian before using in severely debilitated animals and for assistance in the diagnosis, treatment and control of parasitism. Do not mix in feeds containing bentonite.

Store at or below 25° C (77° F).

#### TOP-DRESS DIRECTIONS

LB BODY WEIGHT	OUNCES PER DAY OF PYRANTELCARE™ DAILY DEWORMER 2.11%
250	0.5
500	1.0
750	1.5
1000	2.0
1250	2.5

1 oz. Measuring Scoop Enclosed = 1 Level Scoop per 500 lb. body weight



Manufactured for:  
**FARNAM COMPANIES, INC.**  
P.O. Box 34820 • Phoenix, Arizona 85067-4820  
farnam.com

In case of emergency or for product use information, call 800-234-2269  
Item: 100521750 • RM: 99.0002190.00 • Rev 03-2019

©2019 Farnam Companies, Inc.  
Farnam with design, PyrantelCare and the red to yellow gradient are trademarks of Farnam Companies, Inc.  
Central Garden & Pet and Central Garden & Pet with design are trademarks of Central Garden & Pet Company.

**WARNING:**  
Do not use in horses  
intended for human  
consumption

TAKE TIME  
  
OBSERVE LABEL  
DIRECTIONS

**CENTRAL**  
Garden & Pet



Lot:  
Exp: