Feed for ONE (1) DAY as above. Prior withholding of food or water is not necessary. Retreatment may be required due to the possibility of reinfection. Consult your veterinarian for assistance with the diagnosis and treatment of internal parasites.

#### ANIMAL SAFETY:

There are no known contraindications for the use of fenbendazole in horses. Side effects associated with fenbendazole could not be established in well-controlled safety studies in horses with single doses as high as 454 mg/lb (1,000 mg/kg) and 15 consecutive daily doses of 22.7 mg/lb (50 mg/kg).

Farnam® FenCare™ (fenbendazole) has been evaluated for safety in pregnant mares during all stages of gestation with doses as high as 11.4 mg/lb (25 mg/kg) and in stallions with doses as high as 11.4 mg/lb (25 mg/kg). No adverse effects on reproduction were detected.

# WARNING:

# KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN. NOT FOR USE IN HUMANS.

Do not use in horses intended for human consumption. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report adverse effects, obtain an SDS, or for assistance, contact customer service at 800-234-2269.

#### Manufactured for:

Farnam Horse Products
A Division of Farnam Companies, Inc.
PO Box 34820 • Phoenix, AZ 85067-4820

#### Farnam.com

© 2020 Farnam Companies, Inc.

Item: 100532874 RM: 99.0004145.00

20-10252

Farnam, Farnam with design, and red to yellow color gradation are trademarks of Farnam Companies, Inc.

Store at or below room temperature





Safe-Guard® (fenbendazole) **1.96%**TYPE B MEDICATED FEED

**EQUINE DEWORMER** 



Pelleted Form Easy to Top-Dress on Feed



For the Control of Small and Large Strongyles, Pinworms, and Ascarids



1 Packet Treats up to 1,250 lb Body Weight

# For use in horses only.

Approved by FDA under NADA 131-675
Safe-Guard is a registered trademark of Intervet, Inc., d/b/a Merck Animal Health, a subsidiary of Merck & Co. Inc. All rights reserved.

NET WT. 5 oz (141.7 g)

# INDICATIONS:

For the control of large strongyles (Strongylus edentatus, S. equinus, S. vulgaris, Triodontophorus spp.), small strongyles (Cyathostomum spp., Cylicocyclus spp., Cylicostephanus spp.), pinworms (Oxyuris equi) and ascarids (Parascaris equorum).

### ACTIVE DRUG INGREDIENT:

Fenbendazole 1.96% (556 mg/oz)

#### **GUARANTEED ANALYSIS:**

Crude Protein (not less than) .								10.0%
Crude Fat (not less than)								3.0%
Crude Fiber (not more than)								10.0%
Calcium (not less than)								6.0%
Calcium (not more than)								8.0%
Phosphorus (not less than)								0.5%
Selenium (not less than)						(	),(	02 ppm
Zinc (not less than)								

#### INGREDIENTS:

Ground Wheat, Calcium Carbonate, Soybean Oil, Salt, Urea Formaldehyde, Kaolin, Calcium Sulfate, Silicon Dioxide, Artificial Flavors, Ground Corn Cob, Propionic Acid (preservative), Sodium Carbonate, Calcium Oxide, Lemon Oil, Ammonium Hydroxide, Benzoic Acid (preservative), Phosphoric Acid, Sorbic Acid (preservative), Propylparaben (preservative), Methylparaben (preservative), Butylated Hydroxyanisole (preservative).

#### MIXING DIRECTIONS:

Mix 1 oz Farnam® FenCare™ equine dewormer (containing 556 mg/oz fenbendazole) with 3 oz of non-medicated feed to make 4 oz of a Type C medicated feed (containing the equivalent of 2270 mg/lb).

#### FEEDING DIRECTIONS:

ONE (1) DAY treatment for large strongyles, small strongyles and pinworms: Recommended dose is 5 mg fenbendazole per kg body weight (2.27 mg fenbendazole per pound). Feed 4 oz Type C medicated feed for each 250 pounds of body weight. Example: a 1,250-pound horse would receive 5 oz Farnam® FenCare™ mixed with 15 oz of non-medicated feed for a total of 20 oz Type C medicated feed.

ONE (1) DAY treatment for ascarids: Recommended dose is 10 mg fenbendazole per kg body weight (4.54 mg fenbendazole per pound). Feed 4 oz Type C medicated feed for each 125 pounds of body weight. Example: a 1,250-pound horse would receive 10 oz Farnam® FenCare™ mixed with 30 oz of non-medicated feed for a total of 40 oz. Type C medicated feed.