

CALVENZA[®] EIV/EHV

Boehringer

Equine Rhinopneumonitis-Influenza Vaccine

Killed Virus

Veterinary Use Only

Indications: For the vaccination of healthy, susceptible horses 6 months of age or older, including pregnant mares, as an aid in the reduction of respiratory diseases caused by equine herpesvirus type 1 (EHV-1) and type 4 (EHV-4), and by equine influenza virus types A2 North American and A2 Eurasian.

Administration and Dosage: Shake well before use. Using aseptic technique, vaccinate horses intramuscularly with a 2 mL dose. Administer a second 2 mL dose intramuscularly in 3-4 weeks using a different injection site. Administer a third 2 mL dose in 3-4 weeks by either the intramuscular or intranasal route.

Precautions: Store at 2-7°C. Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur. Antidote: Epinephrine.

Preservatives: Gentamicin and thimerosal.

Manufactured by: **Boehringer Ingelheim Vetmedica, Inc., St. Joseph, Missouri 64506 U.S.A.**

US Vet. Lic. No. 124

Manufactured for: **Boehringer Ingelheim (Canada) Ltd., Burlington, Ontario L7L 5H4**

2 mL syringe	132612-00
10 Doses/20 mL	132606-01

CVP No.: 1230045.5