



SAFE-GUARD®

(FENBENDAZOLE) 0.5%

Medicated Dewormer



...hits the target right where the worms live!

CATTLE - Dairy and Beef: For the removal and control of: Lungworms: (*Dictyocaulus viviparus*); barberpole worms (*Haemonchus contortus*); brown stomach worms (*Ostertagia ostertagi*); small stomach worms (*Trichostrongylus axei*); hookworms (*Bunostomum phlebotomum*); thread-necked intestinal worms (*Nematodirus helvetianus*); small intestinal worms (*Cooperia punctata*, *Cooperia oncophora*); bankrupt worms (*Trichostrongylus colubriformis*); and nodular worms (*Oesophagostomum radiatum*). **EQUINE:** For the removal and control of large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*, *Triodontophorus* spp.), small strongyles (*Cyathostomum* spp., *Cylicocyclus* spp., *Cylicostephanus* spp.), pinworms (*Oxyuris equi*) and ascarids (*Parascaris equorum*).

SWINE - Growing Pigs, Gilts, Pregnant Sows, and Boars: For the removal and control of: Lungworms: (*Metastrongylus apri*, *Metastrongylus pudendotectus*); Gastrointestinal worms: Adult and larvae (L3, L4 stages, liver, lung, intestinal forms) large roundworms (*Ascaris suum*); nodular worms (*Oesophagostomum dentatum*, *O. quadrispinulatum*); small stomach worms (*Hyostrongylus rubidus*); adult and larvae (L2, L3, L4 stages-intestinal mucosal forms) whipworms (*Trichuris suis*) Kidney worms: adult and larvae (*Stephanurus dentatus*). **GROWING TURKEYS:** For the removal and control of gastrointestinal worms: round worms, adults and larvae (*Ascaridia dissimilis*); cecal worms, adult and larvae (*Heterakis gallinarum*) and an important vector of *Histomonas meleagridis* (Blackhead).

ZOO AND WILDLIFE ANIMALS - Bighorn Sheep: Lung worms (*Protostrongylus* spp.). **Feral Swine:** Kidney worms (*Stephanurus edentatus*) round worms (*Ascaris suum*) and nodular worms (*Oesophagostomum edentatum*). **Ruminants (Subfamilies antilopinae, hippotraginae, and caprinae):** Small stomach worm (*Trichostrongylus* spp.); thread necked intestinal worm (*Nematodirus* spp.); barberpole worm (*Haemonchus* spp.) and whipworm (*Trichuris* spp.).

ACTIVE DRUG INGREDIENT

Fenbendazole.....2.27 grams per pound

GUARANTEED ANALYSIS

Crude Protein min..... 12.00%
 Crude Fat min..... 3.00%
 Crude Fiber max..... 16.00%

INGREDIENTS

Processed grain by-products, forage products, roughage products, grain products, calcium carbonate, fat product feed grade, salt, propionic acid, ammonium hydroxide, acetic acid, benzoic acid, sorbic acid, tartaric acid, natural and artificial flavors.

1 cup = 8 oz.

FEEDING DIRECTIONS

CATTLE - Dairy and Beef: 5 mg active drug ingredient Fenbendazole per kg body weight in a ONE (1) DAY treatment (2.27 mg Fenbendazole per pound of bodyweight). **Example:** 1000 lb. cattle should receive 1 lb. of Safe-Guard® 0.5% in one (1) day. No prior withdrawal of feed or water necessary. Under conditions of continued exposure to parasites, retreatment may be needed after 4-6 weeks.

EQUINE: For large strongyles, small strongyles and pinworms, the recommended dose is 5 mg active ingredient Fenbendazole per kg body weight (2.27 mg Fenbendazole per pound) in a ONE (1) DAY treatment. **Example:** a 1000-lb horse should receive 1.0 pounds of Safe-Guard® 0.5% in a ONE (1) DAY treatment. For ascarids, the recommended dose is 10 mg Fenbendazole per kg body weight (4.54 mg fenbendazole per pound) in a ONE (1) DAY treatment. **Example:** a 1000-lb horse should receive 2.0 pounds of Safe-Guard® 0.5% in a ONE (1) DAY treatment. All horses must be eating normally to ensure that each animal consumes an adequate amount of medicated feed. Regular deworming at intervals of 6 to 8 weeks may be required due to the possibility of reinfection. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

SWINE - Growing Pigs, Gilts, Pregnant Sows, and Boars: 9 mg active drug ingredient Fenbendazole per kg body weight (4.08 mg Fenbendazole per pound) to be fed as the sole ration over a period of 3 to 12 days. **Example:** a 200 lb. pig consuming 8 lbs. of feed per day should receive 1.92 ounces of Safe-Guard® 0.5% each day for three (3) days. No prior withdrawal of feed or water necessary.

GROWING TURKEYS: 14.5 grams active drug ingredient Fenbendazole/ ton of feed, to be fed as the sole ration for 6 days. **Example:** 1 lb. of Safe-Guard® 0.5% should be mixed into 313 pounds of feed, to be fed as the sole ration to growing turkeys for 6 consecutive days. No Prior withdrawal of feed or water necessary. Growing turkey feeds containing Safe-Guard® 0.5% can be feed pelleted or as a meal.

ZOO AND WILDLIFE ANIMALS - Big Horn Sheep: 10 mg active drug ingredient Fenbendazole per kg of body weight per day, times (x) 3 days of treatment. **Feral Swine:** 3 mg active drug ingredient Fenbendazole per kg of body weight per day, times (x) 3 days of treatment. **Ruminants (Subfamilies antilopinae, hippotraginae and caprinae)** 2.5 mg active drug ingredient Fenbendazole per kg of body weight per day, times (x) 3 days of treatment. It is recommended that the user exercise judgmental expertise as needed for retreatment within six (6) weeks. This would depend upon the conditions of treated animals, and ambient temperatures.

WARNINGS: KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN. NOT FOR USE IN HUMANS. The Material Safety Data Sheet (MSDS) contains more detailed occupational safety information. To report adverse effects, obtain an MSDS or for assistance, contact customer service at 1-800-441-8272.

Dairy and Beef - Residue warning - Cattle must not be slaughtered within 13 days following last treatment. There are no known contradictions to the use of the drug in cattle. For dairy cattle, there is no milk withdrawal period. **Equine -** Do not use in horses intended for food. **Swine -** Residue warning - There is no pre-slaughter withdrawal period as Safe-Guard® 0.5% can be fed to day of slaughter. **Growing Turkeys -** Residue warning - There is no pre-slaughter withdrawal period as Safe-Guard® 0.5% can be fed to day of slaughter. **Zoo and Wildlife Animals -** Residue warning - Do not use 14 days before or during the hunting season. Animal Safety: No contradictions for the use of Fenbendazole in a zoo environment have been established. Administration to breeding and pregnant ruminants at 2 to 22 times the recommended dose has had no apparent adverse effect.

MANUFACTURED FOR: MARSYT, INC. • ELIZABETHTOWN, PA 17022 • 888-662-7798

NET WEIGHT 5 LBS (2.27 KG)