

**Safe-Guard®** DEWORMER FOR BEEF & DAIRY CATTLE AND GOATS SUSPENSION 10%  
 Intervet/Merck Animal Health  
 (fenbendazole)  
 (100 mg/mL)

**IMPORTANT RESIDUE WARNINGS:**

**Cattle:** must not be slaughtered within 8 days following treatment. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Goats must not be slaughtered for food within 6 days following treatment. Because a withdrawal time in milk has not been established, do not use in lactating goats. For dairy cattle, there is no milk withdrawal period.

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

Keep this and all medication out of the reach of children.

**INDICATIONS:**

**Beef and Dairy Cattle** - 2.3 mg/lb (5 mg/kg) body weight for the removal and control of:

- Lungworm (*Dictyocaulus viviparus*);
- Stomach worm (adults): brown stomach worm (*Ostertagia ostertagi*);
- Stomach worms (adults and fourth stage larvae): barberpole worm (*Haemonchus contortus* and *H. placei*) and small stomach worm (*Trichostrongylus axei*);
- Intestinal worms (adults and fourth stage larvae): hookworm (*Bunostomum phlebotomum*), thread-necked intestinal worm (*Nematodirus helvetianus*), small intestinal worm (*Cooperia punctata* and *C. oncophora*), bankrupt worm (*Trichostrongylus colubriformis*), and nodular worm (*Oesophagostomum radiatum*).

**Goats** - 2.3 mg/lb (5 mg/kg) body weight for the removal and control of:

- Stomach worms (adults): *Haemonchus contortus* and *Teladorsagia circumcincta*.

**DIRECTIONS:** Determine the proper dose according to estimated body weight. Administer orally. The recommended dose of 2.3 mg/lb (5 mg/kg) of body weight is achieved when 2.3 mL of the drug are given for each 100 lbs body weight.

**EXAMPLES:**

Cattle:

Goats:

Body Weight	Amount
100 lb	2.3 mL
200 lb	4.6 mL
300 lb	6.9 mL
400 lb	9.2 mL
500 lb	11.5 mL
1,000 lb	23.0 mL
1,500 lb	34.5 mL

Body Weight	Amount
25 lb	0.6 mL
50 lb	1.2 mL
75 lb	1.7 mL
100 lb	2.3 mL
125 lb	2.9 mL

Under conditions of continued exposure to parasites, retreatment may be needed after 4 to 6 weeks.

Store at or below 25°C (77°F). Protect from freezing. Shake well before use.

Restricted drug (California) - use only as directed.  
 For Use in Animals Only.

Made in France  
 Distributed by: Intervet Inc (d/b/a Merck Animal Health), Madison, NJ 07940

NADA # 128-620, Approved by FDA

	Code No.	
1,000 mL (33.8 fl oz)	069293	Rev. 06/14 189850 R1 127879 R2
1 Gallon (3,785 mL)		153593 R2 151653 R2
10 Liter		09/2015 159156 R2 158338 R2