

# KOPERTOX®

## Topical Antifungal

### Active Ingredient(s):

Copper naphthenate 37.5%

Inert ingredients 62.5%

Total 100.0%

**Indications:** Recommended as an aid in treating horses and ponies with thrush due to organisms susceptible to copper naphthenate.

### Dosage and Administration:

**General directions:** Clean the hoof thoroughly removing debris and necrotic material prior to application of KOPERTOX®. Apply daily to affected hoofs with a narrow paint brush (about 1") until fully healed.

**Note:** KOPERTOX® is easily removed from hands, clothing and surfaces with light grade fuel oil or any type of lighter fluid.

**Precaution(s):** Store at controlled room temperature 15°C to 30°C (59° to 86°F).

**Caution(s):** Combustible mixture. Use in a well-ventilated place. Avoid fire, flame, sparks or heaters.

If swallowed, do not induce vomiting; call physician immediately. Avoid breathing vapor. Avoid contact with skin and eyes.

Do not allow runoff of excess KOPERTOX® onto hair since contact with KOPERTOX® may cause some hair loss. Do not contaminate feed.

Keep out of reach of children and pets.

For external use only.

**Warning(s):** Do not use on animals which are raised for food production.

Not for use on horses intended for food.

**Presentation:** ½ pint (8 oz., 236 mL) (NDC 0856-9910-01) and 1 pint (16 oz., 473 mL) (NDC 0856-9910-02), 12 in a case.