Indications: For vaccination of healthy horses 4 months of age or older as an aid in the prevention of disease caused by Eastern and Western Encephalomyelitis, and tetanus; as an aid and an aid in the reduction of severity of clinical disease caused by West Nile Virus; as an aid in the reduction of respiratory types 1 and 4 (EHV-1 and EHV-4); and as clade 1 (OH/03), and Florida sublineage at least 6 months against OH/03. The

Administration and dosage: Shake well before use. Using aseptic technique, inoculate horses intramuscularly with a 1 ml dose. Administer a second 1 mL dose intramuscularly in 3-4 weeks using a different injection site. Revaccinate annually and prior to anticipated exposure with a single 1 ml Intramuscular dose.

Encephalomyelitis-Rhinopneumonitis-Influenza-West Nile Virus Vaccine Eastern & Western, Killed Virus Tetanus Toxoid Vetera © Gold<sup>XP</sup> Precautions: Store at 35-45°F (2-7°C). Protect from freezing, Do not vaccinate within 21 days before slaughter. Use entire contents when first opened. Anaphylactoid reactions may occur following use. Antidote: Epinephrine.

This product has been tested under laboratory conditions and shown to meet all Federal standards for safety and efficacy in normal, healthy, immunocompetent animals. This level of performance may be affected by conditions such as stress, weather, nutrition, disease, parasitism, other treatments, individual lidosyncrasies, or impaired immunological competency. These factors should be considered by the user when evaluating product performance or freedom from reactions.

Preservatives: Gentamicin and Amphotericin B.

## For Veterinary Use Only

Boehringer Ingelheim Vetmedica GMBH.
© 2013 Boehringer Ingelheim

© 2013 Boehringer Ingelheim Vetmedica, Inc. All Rights Reserved.

Boehringer Ingelheim Vetmedica, Inc. St. Joseph, MO 64506 US Vet. Lic. No. 124 US Patent No. 7572620. Used under license. Code 137-125

