Clovite® Conditioner (ZOETIS INC.)

Label Use/Dose

ZOETIS INC.

333 PORTAGE STREET, KALAMAZOO, MI, 49007

Telephone: 269-833-4000

Customer Service: 800-733-5500 and 800-793-0596

Veterinary Medical Investigations & Product Support: 800-366-5288

Technical Services (USA): 800-366-5288

Website: www.zoetis.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

CLOVITE® CONDITIONER

Zoetis

VITAMIN SUPPLEMENT FOR ALL SPECIES OF ANIMALS

MINIMUM VITAMIN GUARANTEE

(each pound contains)

VITAMIN A	110,000 IU (242 IU/g)
VITAMIN D	50,000 IU (110 IU/g)
VITAMIN B ₁₂	136 mcg (0.3 mcg/g)

INGREDIENTS: Soybean meal, Soy flour, Vitamin A & D oil, Dicalcium phosphate, Vitamin B₁₂ supplement, Vitamin A & D₃ supplement, Choline chloride, Vitamin E supplement, Calcium pantothenate, Niacin, Pyridoxine hydrochloride, Thiamine hydrochloride, Riboflavin, Biotin.

SUGGESTED USAGE:

Young Foals and Weanlings: 1 to 2 tablespoonfuls daily.

Brood Mares: (latter half of pregnancy and lactation) 2 tablespoonfuls twice a day.

Ponies: 1 tablespoonful daily.

Colts, Stallions and Horses in Training: 1 tablespoonful per 400 lb daily.

All Farm Animals: 1 to 3 lb to each 100 lb feed.

Mature Cattle: (milking or on full feed) 1 to 3 lb to each 100 lb feed. Many dairymen prefer adding a handful to each feeding. Feedlot operators simply top off feed with the prescribed amount of Clovite.

Calves: (bucket fed) 1 to 2 tablespoonfuls daily in milk or ground feed.

Range Cattle: Supplemental feeding is simplified by mixing with salt - 1 part to 10 parts salt, or 2 1/2 lb to 25 lb salt. (These calculations are based on normal consumption of 1/4 lb of salt per day by a 1,000 lb cow and 1/8 lb per day by a 400 lb steer.) To prevent the oxidation that tends to destroy all Vitamin A compounds on prolonged exposure, mix only the amount normally consumed in a 3-week period.

Poultry: 1 lb to each 100 lb feed.

Baby Pigs: 3% of the creep-feed.

Dogs and Cats: 1 level teaspoonful per each 10 lb of body weight daily. This dosage should be doubled for growing pups and kittens, lactating animals and animals in the last half of pregnancy. Dosage may be varied as prescribed by the veterinarian.

Mink: 1 level teaspoonful daily.

Fort Dodge Animal Health, Fort Dodge, Iowa 50501 USA

NET WT	
5 LB (2.2 KG)	00642
	2336I
25 LB	2331M

NAC No.: 10030344