## EQUINE POTOMAVAC™

Merial

Ehrlichia Risticii Bacterin

PHF

DESCRIPTION: EQUINE POTOMAVAC™ is a liquid suspension of inactivated Ehrlichia risticii. Safety and efficacy of this product have been demonstrated in controlled vaccination challenge tests and in thorough field evaluations.

INDICATIONS: Recommended for the vaccination of healthy horses as an aid in the prevention of Potomac Horse Fever (equine monocytic ehrlichiosis) caused by E. risticii.

DOSAGE: For primary vaccination, aseptically inject 1 mL (1 dose) intramuscularly into healthy horses 3 months of age or older. Revaccinate with a second 1 mL dose 3 to 4 weeks later. Revaccinate annually with a single 1 mL dose.

PRECAUTIONS: Store at 2-7°C (35-45°F). Do not freeze. Use entire contents when first opened. Shake well before using. Contains gentamicin as a preservative. Do not vaccinate within 21 days prior to slaughter. In rare instances, administration of vaccines may cause lethargy, fever, and inflammatory or hypersensitivity types of reactions. Treatment may include antihistamines, anti-inflammatories, and/or epinephrine. Vaccinating animals whose immune response is compromised by stress, disease, etc. may not produce the desired results.

SOLD TO VETERINARIANS ONLY.

For Veterinary Use Only.

™POTOMAVAC is a trademark of MERIAL.

Manufactured By MERIAL INC., Athens, GA 30601 USA

U.S. Vet. Lic. No. 298

1-888-Merial-1 (1-888-637-4251)

Distributed In Canada By MERIAL CANADA, INC., Baie d'Urfé, Qc, H9X 3V1