

Drug Facts

Active ingredient (in each 1 tsp)

Pyrantel base: 250 mg.....Pinworm treatment
(as Pyrantel Pamoate)

Purpose

Indication For the treatment of pinworms.

Warnings

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center immediately.

If you are pregnant or have liver disease, do not take this product unless directed by a physician.

When using this product, abdominal cramps, nausea, vomiting, diarrhea, headache, or dizziness sometimes occur after taking this drug. If any of these conditions persist, consult a physician.

Directions

- Shake well before use.
- Adults, children 12 years of age and over, and children 2 years to under 12 years of age: Oral dosage is a single dose of 5 milligrams of pyrantel base per pound, or 11 milligrams per kilogram, of body weight not to exceed 1 gram (4 teaspoonfuls).

WEIGHT	DOSAGE (TAKEN AS A SINGLE DOSE)
Less than 25 lbs or under 2 years old	Do not use unless directed by a physician
25 to 37 lbs	½ teaspoonful (125 mg)
38 to 62 lbs	1 teaspoonful (250 mg)
63 to 87 lbs	1½ teaspoonfuls (375 mg)
88 to 112 lbs	2 teaspoonfuls (500 mg)
113 to 137 lbs	2½ teaspoonfuls (625 mg)
138 to 162 lbs	3 teaspoonfuls (750 mg)
163 to 187 lbs	3 ½ teaspoonfuls (875 mg)
188 lbs and over	4 teaspoonfuls (1,000 mg)

- Dosage information: Each 5 mL (1 teaspoonful) of Pyrantel Pamoate Suspension contains the equivalent of 250 mg of pyrantel base.
- Read package insert carefully before taking this medication.

Directions continued ►

NDC: 62337-415-32



PYRANTEL PAMOATE SUSPENSION

(Pyrantel base: 50 mg / mL)

PINWORM TREATMENT

 **COLOMBIA**
LABORATORIES™
944 Nandino Blvd.
Lexington, KY 40511

Net Contents: 32 fl oz (946.4 mL)

Drug Facts (continued)

Directions (continued)

- Take only according to directions and do not exceed the recommended dosage unless directed by a physician.
- Medication should only be taken one time as a single dose; do not repeat treatment unless directed by a physician.
- When one individual in a household has pinworms, the entire household should be treated unless otherwise advised. See Warnings.
- If any worms other than pinworms are present before or after treatment, consult a physician.
- If any symptoms or pinworms are still present after treatment, consult a physician.
- This product can be taken any time of day, with or without meals. It may be taken alone or with milk or fruit juice.
- Use of a laxative is not necessary prior to, during, or after medication.

Other information

- Recommended storage: Store at temperature between 15°C – 30°C (59°F – 86°F).

Inactive ingredients

Citric Acid, Potassium Sorbate, Purified Water, USP, Sodium Benzoate, Sucrose, Vanilla Flavor, and Xanthan Gum

Questions? Call 859-254-1221 or toll-free 800-621-8829.

DO NOT USE IF SEAL UNDER CAP IS BROKEN OR MISSING.

Item No. 09073



L6366-0519