

Liquid Wormer 2X

(pyrantel pamoate)

Canine Anthelmintic Suspension

DIRECTIONS FOR USE:

For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8 and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent T. canis reinfestation. Administer one full teaspoonful (5 mL) for each 10 lbs. of body weight.

For the removal of large roundworms (ascarids) and hookworms. Administer one full teaspoonful (5 mL) for each 10 lbs. of body weight. It is not necessary to withhold food prior to treatment.

Dogs usually find this wormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption.

It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2 to 4 weeks after first treatment.

RECOMMENDED STORAGE:
STORE BELOW 30°C (86°F)

Read Accompanying Professional Information

NDC 30798-760-22

EACH 2 OZ BOTTLE TREATS 120 LBS

Liquid Wormer 2X

(pyrantel pamoate)

5 ML PER 10 LB BODY WEIGHT

Canine Anthelmintic Suspension

Removes
Large Roundworms
& Hookworms
in Puppies
& Dogs

durvet



MADE IN USA

Active Ingredient

4.54 mg
of Pyrantel
Base as
Pyrantel
Pamoate
per mL

FOR ANIMAL USE ONLY
KEEP OUT OF REACH OF CHILDREN
ANADA 200-007 APPROVED BY FDA
NET CONTENTS: 2 FL OZ (60 mL)

Liquid Wormer 2X

(pyrantel pamoate)

INDICATIONS FOR USE:

To prevent reinfestation of Toxocara canis in puppies and adult dogs and in lactating bitches after whelping.

For the removal of large roundworms, (Toxocara canis and Toxascaris leonina) and hookworms, (Ancylostoma caninum and Uncinaria stenocephala) in dogs and puppies.

The presence of these parasites should be confirmed by laboratory fecal examination. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

TAKE TIME



OBSERVE LABEL
DIRECTIONS

IMPORTANT: SHAKE WELL BEFORE USE

**WARNING: KEEP OUT
OF THE REACH OF CHILDREN**

durvet

www.durvet.com

LOT & EXP IMPRINT COPY

INDICATIONS FOR USE

To prevent reinfestation of (Toxocara canis) in puppies and adult dogs and in lactating bitches after whelping.

For the removal of large roundworms, (Toxocara canis and Toxascaris leonina) and hookworms, (Ancylostoma caninum and Uncinaria stenocephala) in dogs and puppies.

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

WARNING: KEEP OUT OF REACH OF CHILDREN
Read Accompanying Professional Information

Recommended Storage: Store Below 30°C (86°F)



www.durvet.com

Manufactured for Durvet, Inc.
100 S.E. Magellan Drive,
Blue Springs, Missouri 64014
ISS12XB04

NDC 30798-760-22

durvet

Liquid Wormer

(pyrantel pamoate) **2X**

Canine Anthelmintic Suspension

Active Ingredient: 4.54 mg of
pyrantel base as pyrantel pamoate per mL

FOR ANIMAL USE ONLY

ANADA 200-007 APPROVED BY FDA
NET CONTENTS: 2 FL OZ (60 ML)

IMPORTANT: SHAKE WELL BEFORE USING

DIRECTIONS FOR USE: For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8 and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent T. canis reinfestation. Administer one full teaspoonful (5 mL) for each 10 lbs. of body weight.

For the removal of large roundworms (ascarids) and hookworms. Administer one full teaspoonful (5 mL) for each 10 lbs. of body weight. It is not necessary to withhold food prior to treatment. Dogs usually find this wormer very palatable and will lick the dose from the bowl willingly. If there is a reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. The presence of large roundworms (ascarids) and hookworms should be confirmed by lab fecal examination.

It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2 to 4 weeks after first treatment.

Take Time



**Observe Label
Directions**

IMPRINT AREA